

NCLEX RECALL BINDER

HIGH-YIELD · VISUAL · EASY TO REMEMBER



How to use this binder

mnemonurse.com/recall-binder

Meet the characters

158 of the hardest, highest-yield NCLEX topics turned into 158 patient characters you actually remember — with signs, nursing pearls, lab values, flashcards and rationale-backed practice questions. Every hard topic gets a face, a story hook and the exact findings you will be tested on.

How each character is laid out

Every character gets four pages. Page one is the portrait page: the character, the memory hook, why this character matters, and the story arc in four beats — the setup, the turn, the crisis, the save. Page two explains what goes wrong, what normal looks like, and a closer look at the findings that matter. Page three collects the warning signs and the priority action. Page four gives you a practice question or a rapid-recall check, the “remember it” teaching, and the memory line that locks the character in.

Study tools at the back

Behind the characters you will find a gallery of all 158 characters, the Rapid-Recall Flashcard Deck (cover the answer, say it out loud, then check yourself), a day-by-day study planner, and four quick indexes: an A-Z character directory, a memory-line index, a warning-sign quick index, and a priority-action quick index.

Pace yourself

Pace options: Steady = 2 characters a day, Standard = 3 a day, Cram = 6 a day. The planner at the back is set to the Standard pace. Tick each box, then re-test yourself the next morning.

Table of Contents

Tap or click any chapter to jump straight to it.

Characters

1. Acidotic Andy - Metabolic Acidosis	9
2. Alkalotic Alice - Metabolic Alkalosis	13
3. Retaining Ray - Respiratory Acidosis	17
4. Panicking Pam - Respiratory Alkalosis	21
5. Potassium Pete - Hyperkalemia	25
6. Low-K Lena - Hypokalemia	29
7. Salty Sal - Hypernatremia	33
8. Diluted Dana - Hyponatremia	37
9. Chvostek Chuck - Hypocalcemia	41
10. Stony Stella - Hypercalcemia	45
11. Magnesium Maggie - Magnesium Imbalance	49
12. Septic Sam - Distributive Shock	53
13. Pumpless Paul - Cardiogenic Shock	57
14. Bleeding Bea - Hypovolemic Shock	61
15. Anaphylactic Ana - Anaphylaxis	65
16. Embolic Emma - Pulmonary Embolism	69
17. Rigid Rita - ARDS	73
18. Wheezing Wanda - Asthma Exacerbation	77
19. Consolidated Carl - Pneumonia	81
20. Vented Vince - Mechanical Ventilation	85
21. Ketone Kyle - Diabetic Ketoacidosis	89
22. Hyperosmolar Hank - Hyperosmolar Hyperglycemic State	93
23. Shaky Shawna - Hypoglycemia	97
24. Storming Tori - Thyroid Storm	101
25. Sluggish Myra - Myxedema Coma	105
26. Addison Ada - Adrenal Crisis	109
27. Cushing Cora - Cushing's Syndrome	113

Table of Contents — continued

28. Dry Dwayne - Diabetes Insipidus	117
29. Pressured Priya - Increased Intracranial Pressure	121
30. Stroke Stan - Ischemic Stroke	125
31. Seizing Simone - Seizures	129
32. Failing Frank - Heart Failure	133
33. Infarcting Ivan - Myocardial Infarction	137
34. Fibrillating Fay - Atrial Fibrillation	141
35. Warfarin Wally - Anticoagulation	145
36. Ammonia Amy - Cirrhosis	149
37. Anuric Alan - Acute Kidney Injury	153
38. Digoxin Dot - Digoxin Toxicity	157
39. Preeclamptic Penny - Preeclampsia	161
40. Pancreatic Pablo - Acute Pancreatitis	165
41. Sober Sully - Alcohol Withdrawal	169
42. Manic Mia - Bipolar Mania	173
43. Voices Victor - Schizophrenia	177
44. Sinking Sadie - Major Depression	181
45. Flashback Felix - PTSD	185
46. Narcan Nate - Opioid Overdose & Withdrawal	189
47. Mirror Mila - Anorexia Nervosa & Refeeding Syndrome	193
48. Serotonin Sylvia - Serotonin Syndrome	197
49. Wheezy Winnie - Bronchiolitis	201
50. Sickled Simon - Sickle Cell Vaso-Occlusive Crisis	205
51. Cystic Cassie - Cystic Fibrosis	209
52. Strawberry Kai - Kawasaki Disease	213
53. Telescope Tommy - Intussusception	217
54. Droopy Dani - Infant Dehydration & Gastroenteritis	221
55. Lead-ed Liam - Lead Poisoning	225
56. Gushing Gwen - Postpartum Hemorrhage	229
57. Previa Paula - Placenta Previa	233
58. Abrupting Abby - Placental Abruption	237
59. Sugary Selena - Gestational Diabetes	241

Table of Contents — continued

60. Decel Dina - Fetal Heart Rate Decelerations	245
61. Yellow Yusuf - Newborn Hyperbilirubinemia & Phototherapy	249
62. Cord Callie - Umbilical Cord Prolapse	253
63. Feverish Fiona - Postpartum Infection & Mastitis	257
64. Neutropenic Nia - Neutropenia & Chemotherapy Precautions	261
65. Lysing Luis - Tumor Lysis Syndrome	265
66. Transfusing Tina - Blood Transfusion Reactions	269
67. Clotless Clara - Disseminated Intravascular Coagulation	273
68. Pale Priscilla - Iron Deficiency Anemia	277
69. Spore Sean - Clostridioides difficile	281
70. CD4 Hana - HIV & Opportunistic Infections	285
71. Butterfly Bella - Systemic Lupus Erythematosus	289
72. Compartment Cody - Fractures & Compartment Syndrome	293
73. Reflexive Rex - Spinal Cord Injury	297
74. Myasthenic Marge - Myasthenia Gravis	301
75. Tremor Tomas - Parkinson's Disease	305
76. Barrel Bart - COPD	309
77. Tarry Tara - Upper GI Bleed	313
78. Blocked Bruno - Bowel Obstruction	317
79. Crisis Camila - Hypertensive Crisis	321
80. Aneurysm Arturo - Abdominal Aortic Aneurysm	325
81. Claudicating Clara - Peripheral Arterial Disease vs Venous Insufficiency	329
82. Endocarditis Eli - Infective Endocarditis	333
83. Pericardial Pearl - Pericarditis	337
84. Valve Vera - Valvular Heart Disease	341
85. Paced Pia - Pacemakers	345
86. Clotted Cliff - Deep Vein Thrombosis	349
87. Appendix Annie - Appendicitis	353
88. Gallstone Gina - Cholecystitis & Cholelithiasis	357
89. Crohn's Colin - Inflammatory Bowel Disease	361
90. Diverticular Dee - Diverticulitis	365
91. Hepatitis Hugo - Viral Hepatitis A, B & C	369

Table of Contents — continued

92. Dialysis Dina - Chronic Kidney Disease	373
93. Kidney Kwame - UTI & Pyelonephritis	377
94. Prostate Percy - BPH & TURP Care	381
95. Burned Bella - Burns	385
96. Ulcer Uma - Pressure Injury Staging & Prevention	389
97. Confused Connie - Delirium vs Dementia & Restraint Alternatives	393
98. Delegating Deb - Delegation	397
99. Insulin Ike - Insulin	401
100. Triage Theo - Disaster Triage	405
101. Consent Carmen - Informed Consent, Advance Directives & Privacy	409
102. ACE Angie - ACE Inhibitors	413
103. Beta Betty - Beta Blockers	417
104. Lasix Larry - Loop Diuretics	421
105. Lithium Lila - Lithium	425
106. Tyramine Tessa - MAOIs & the Tyramine Hypertensive Crisis	429
107. Nitro Nina - Nitroglycerin: Sublingual Rules, Headache & Dangerous Combinations	433
108. Heparin Holly - Heparin	437
109. Phenytoin Phil - Phenytoin: Narrow Range, Nystagmus & Gummy Smiles	441
110. Rhabdo Rory - Rhabdomyolysis: Tea-Colored Urine, Potassium & Kidney Rescue	445
111. Osteo Opal - Osteoporosis: Silent Bone Loss, Fall Prevention & Bisphosphonates	449
112. Glaucoma Gloria - Glaucoma: Silent Peripheral Loss vs Acute Angle-Closure Emergency	453
113. Meningitis Mia - Bacterial Meningitis	457
114. Sclerosis Sadie - Multiple Sclerosis: Demyelination, Relapses & Heat Intolerance	461
115. Gout Gus - Gout: Uric Acid Crystals, the Angry Big Toe & Purine Teaching	465
116. Meniere Mira - Meniere's Disease: Vertigo, Tinnitus & Hearing Loss	469

Table of Contents — continued

117. Rheumatoid Rosa - Rheumatoid Arthritis: Symmetric Joints, Morning Stiffness & DMARDs	473
118. Obsessive Owen - OCD: Obsessions, Compulsions & Anxiety Relief	477
119. Hyperemesis Hazel - Hyperemesis Gravidarum: Vomiting, Ketones & Dehydration	481
120. Mastitis Mabel - Mastitis: Flu-Like Fever, a Red Wedge & Keep Nursing	485
121. Nephrotic Nolan - Childhood Nephrotic Syndrome: Puffy Eyes, Frothy Urine & Protein Loss	489
122. Phosphate Franklin - Phosphorus Imbalance	493
123. Overloaded Olive - Fluid Volume Overload	497
124. Collapsing Callum - Tension Pneumothorax	501
125. Ascending Abe - Guillain-Barré Syndrome	505
126. Sundowning Sid - Alzheimer's Disease	509
127. Quarantined Quinn - Tuberculosis	513
128. Croupy Cora - Croup	517
129. Fractured Frida - Hip Fracture	521
130. Shingles Sheila - Herpes Zoster	525
131. Warfarin Winnie - Warfarin	529
132. Slow-Drip Delia - IV Potassium Safety	533
133. Central Line Cesar - Central Lines	537
134. Infiltrated Ivy - Infiltration	541
135. Trach Tomas - Tracheostomy	545
136. Feeding Fiona - Enteral Feeding	549
137. Steroid Stan - Corticosteroid Therapy	553
138. Levo Lucy - Levothyroxine: Empty Stomach, Same Time, Lifelong — and the Signs of Too Much	557
139. Vanco Vera - Vancomycin: Slow the Infusion, Watch the Trough, Protect the Kidneys and Ears	561
140. Oxygen Omar - Oxygen Delivery Devices: Matching the Mask to the Need — and Fire Safety	565
141. Post-Op Priya - Postoperative Complications: The Day-by-Day Timeline Every Nurse Watches	569

Table of Contents — continued

142. PCA Pablo - PCA & Opioid Safety: Only the Patient Presses the Button	573
143. Restraint Rita - Restraints & Fall Prevention: Least Restrictive First, and the Order Rules	577
144. Vaccine Vivian - Immunizations: Live Vaccines, Real Contraindications and the Reactions to Expect	581
145. TPN Tessa - Total Parenteral Nutrition: High Sugar, Central Line, and Never Stop It Suddenly	585
146. Listening Theo - Therapeutic Communication: What to Say, and What Never to Say	589
147. Milestone Maya - Growth, Development and Age-Appropriate Safety	593
148. Labor Lena - Stages of Labor	597
149. Screening Sam - Cancer Warning Signs	601
150. Pressure Pam - Antihypertensive Classes	605
151. Ectopic Erica - Ectopic Pregnancy	609
152. Tet Spell Tilda - Tetralogy of Fallot	613
153. Pyloric Piper - Pyloric Stenosis	617
154. Phantom Petra - Amputation Care	621
155. Glomerular Greta - Glomerulonephritis	625
156. B12 Bianca - Pernicious Anemia	629
157. Retinal Rhea - Retinal Detachment	633
158. Ostomy Oscar - Ostomy Care	637

Reference

Meet the 158 Recall Characters	641
Rapid-Recall Flashcard Deck	668
Your Study Planner	734
Character Directory (A-Z)	742
Memory-Line Index	745
Warning-Sign Quick Index	750
Priority-Action Quick Index	759

1. Acidotic Andy

Metabolic Acidosis



Kussmaul breathing, fruity breath, pH ↓ · HCO_3^- ↓

Andy pulls into the DKA Diner running on empty, orders a fruity milkshake and starts breathing like a steam engine. Every deep breath is his lungs trying to blow the acid out of a body that has already run out of insulin.

WHY THIS CHARACTER MATTERS

Andy's body is drowning in acid. Look for deep, rapid Kussmaul respirations blowing off CO_2 , confusion, warm flushed skin and hyperkalemia. Labs: pH < 7.35, HCO_3^- < 22, anion gap > 12. Treat the cause first — insulin plus fluids for DKA, not reflexive bicarb.

The story arc

1. The setup

Andy skips two days of insulin during finals week and lives on soda. Cells starve while sugar piles up in the blood outside them.

2. The turn

By morning he is breathing like a steam engine — deep, tireless, loud — and his roommate smells nail polish on his breath.

3. The crisis

In triage his pH reads 7.12, his anion gap is 22, and his potassium comes back high even though he has been urinating out potassium for two days.

4. The save

The nurse hangs isotonic fluids, checks potassium before touching the insulin drip, then titrates and rechecks — no bicarb hero moves.

1. Acidotic Andy • What goes wrong

Metabolic Acidosis



What goes wrong

Two days of missed insulin plus sugary sodas starve his cells while sugar piles up outside them; osmotic diuresis drains water and electrolytes, and ketone acids overwhelm the buffers into high anion gap metabolic acidosis — his lungs blow off carbon dioxide in compensation, but they cannot outpace the acid. By morning he is slumped at his desk, obtunded, breathing like a steam engine.

NORMAL — THE BASELINE

On his basal insulin, Andy's glucose enters his cells, no ketones form, and his bicarbonate buffer idles comfortably with pH in range. Finals week comes and goes without incident.

Closer look

Acid load — With two days of insulin skipped, Andy's liver metabolizes fatty acids into an avalanche of acetoacetic and beta-hydroxybutyric acids. His anion gap spikes to 22 mEq/L — the diagnostic signature: $\text{pH} < 7.35$, $\text{HCO}_3^- < 22$, anion gap > 12 .

Bicarbonate buffer — The acid consumes his buffer: serum bicarbonate drops to a dismal 8 mEq/L while arterial pH reads 7.12. The rule reads straight off the glass: when pH is low and bicarbonate is low, the metabolic system is below normal.

Breathing compensation — Every deep breath is his lungs trying to blow the acid out: loud, deep, tireless Kussmaul respirations drive his PaCO_2 down to 24 mmHg in partial compensation. His roommate smells nail polish — acetone — on that breath.

pH gauge — Watch the potassium trick on this gauge's neighbor: serum potassium reads 5.8 mEq/L because acidosis forces potassium out of cells — high in the blood even though total body stores are depleted after two days of urinating it away. Never start the insulin drip until potassium is verified at or above 3.3 mEq/L.

1. Acidotic Andy · Warning signs

Metabolic Acidosis



Warning signs

- pH 7.12 — below 7.35
- Bicarbonate 8 mEq/L — below 22
- Anion gap 22 — above 12
- Deep, rapid, loud Kussmaul respirations
- Fruity acetone breath — smells like nail polish
- Confusion, obtundation — responding only to noxious stimuli
- Warm, flushed skin
- Hyperkalemia — potassium 5.8 mEq/L despite depleted total stores
- Deep muscular weakness and generalized abdominal cramping
- Nausea and relentless thirst
- PaCO₂ 24 mmHg — lungs blowing off carbon dioxide

Priority action

Treat the cause first — insulin plus fluids for DKA, not reflexive bicarb. Hang isotonic fluids to restore volume and renal perfusion, check potassium before touching the insulin drip (at or above 3.3 mEq/L), then titrate the regular insulin infusion with continuous cardiac telemetry, hourly neuro checks, and urine output charting. Intravenous sodium bicarbonate is strictly withheld — rapid alkalization risks rebound hypokalemia and paradoxical central nervous system acidosis.

1. Acidotic Andy · Check yourself

Metabolic Acidosis



Practice question

Andy is admitted in diabetic ketoacidosis with pH 7.22 and bicarbonate 12 mEq/L. Before the insulin infusion is started, his potassium returns at 2.9 mEq/L. What is the nurse's priority action?

- A. Hold the insulin infusion, notify the provider, and anticipate potassium replacement first
- B. Start the insulin infusion immediately, because insulin is the definitive DKA treatment
- C. Start the insulin and request an IV bicarbonate infusion to correct the pH
- D. Start the insulin and recheck the potassium in 4 hours

ANSWER: A

Insulin drives potassium into the cells, so starting it when potassium is already below about 3.3 mEq/L can cause a dangerous further drop and cardiac dysrhythmias; potassium must be replaced first. Although his presenting potassium may look normal or high because acidosis shifts potassium out of cells, total body potassium in DKA is depleted. Reflexive bicarbonate is also not the treatment — fluids and insulin correct the underlying ketoacidosis.

Remember it

To master metabolic acidosis for the NCLEX, picture Acidotic Andy panting like a steam engine at the diner, blowing off carbon dioxide with deep Kussmaul respirations while smelling of fruity acetone. Remember his core diagnostic rule: when pH is low and bicarbonate is low, the metabolic system is below normal. Keep his potassium trick locked in your clinical memory because acidotic patients frequently present with high serum potassium despite depleted total body stores. On exam day, watch for the classic trap: never rush to hang an intravenous bicarbonate infusion, and never start an intravenous regular insulin drip until you verify that serum potassium is at or above 3.3 mEq/L. Fluids and insulin correct the underlying ketoacidosis while protecting the heart from deadly dysrhythmias.

MEMORY LINE

Kussmaul breathing, fruity breath, pH ↓ · HCO₃⁻ ↓

2. Alkalotic Alice

Metabolic Alkalosis



***Vomiting and diuretics, tingling fingers, pH
 $\uparrow$ · HCO_3^- $\uparrow$***

Alice spends three days vomiting after a stomach bug and keeps her water pill going the whole time. Every heave sends hydrochloric acid down the drain until her blood turns sweet and her fingers start to buzz.

WHY THIS CHARACTER MATTERS

Alice loses acid instead of gaining base — protracted vomiting, nasogastric suction or loop diuretics. Expect tingling, tremors, tetany and shallow slow breathing as her lungs try to hold CO_2 . Labs: $\text{pH} > 7.45$, $\text{HCO}_3^- > 26$, K^+ and Cl^- low.

The story arc

1. The setup

Three days of vomiting, no fluids kept down, and a furosemide dose she never skipped.

2. The turn

Her fingers and lips start tingling, her hands cramp when the blood pressure cuff inflates.

3. The crisis

Her breathing slows and shallows out; the monitor shows a run of PVCs while her K^+ reads 2.9.

4. The save

Normal saline with potassium chloride replaces the chloride she vomited away, and the diuretic is held.

2. Alkalotic Alice • What goes wrong

Metabolic Alkalosis



What goes wrong

Three days of vomiting with no fluids kept down, and a furosemide dose she never skipped, drain hydrochloric acid, potassium, and chloride. Her blood turns alkaline, free ionized calcium falls, and tetany and PVCs follow.

NORMAL — THE BASELINE

Stomach acid stays in the stomach, bicarbonate stays in range, and potassium and chloride stay replete — pH balanced.

Closer look

Stomach acid drain — Alice loses acid instead of gaining base: every heave sends hydrochloric acid down the drain — protracted vomiting, nasogastric suction or loop diuretics. She spends three days vomiting after a stomach bug and keeps her water pill (furosemide) going the whole time.

Bicarbonate and pH — Labs: $\text{pH} > 7.45$, $\text{HCO}_3^- > 26$, K^+ and Cl^- low — purging hydrochloric acid leaves the blood elevated in bicarbonate and dangerously low in potassium and chloride.

Tingling fingers and Trousseau spasm — Expect tingling, tremors, tetany: her fingers and lips start tingling and her hands cramp when the blood pressure cuff inflates. An alkaline pH forces calcium to bind tightly to albumin, lowering free ionized calcium and triggering positive Trousseau and Chvostek signs.

Slow breathing — Her breathing slows and shallows out — shallow slow breathing as her lungs try to hold CO_2 ; the body's respiratory attempt to hold onto acidic carbon dioxide.

Potassium and chloride — Her K^+ reads 2.9 and the monitor shows a run of PVCs; the priority treatment is isotonic sodium chloride fluid replacement to restore chloride.

2. Alkalotic Alice • Warning signs

Metabolic Alkalosis



Warning signs

- Three days of vomiting, no fluids kept down, diuretic never skipped
- Fingers and lips tingling
- Hands cramp when the blood pressure cuff inflates (Trousseau)
- Tremors and tetany
- Breathing slow and shallow
- K^+ 2.9 with a run of PVCs on the monitor
- Labs: $pH > 7.45$, $HCO_3^- > 26$, K^+ and Cl^- low

Priority action

Give normal saline with potassium chloride to replace the chloride she vomited away — isotonic sodium chloride fluid replacement to restore chloride is the priority treatment — and hold the diuretic.

2. Alkalotic Alice • Check yourself

Metabolic Alkalosis



Practice question

Alice's hand spasms when the blood pressure cuff inflates. The nurse documents this as:

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it

To remember this condition on exam day, picture Alkalotic Alice holding her stomach while her fingers buzz and twist from tetany after days of vomiting stomach acid and swallowing diuretic water pills. Link the name Alice to the phrase All Loss of Acid, reminding you that purging hydrochloric acid leaves the blood elevated in bicarbonate and dangerously low in potassium and chloride. Watch for her slow, shallow breathing, which represents the body's respiratory attempt to hold onto acidic carbon dioxide. For the NCLEX, remember that metabolic alkalosis frequently masquerades as hypocalcemia because an alkaline pH forces calcium to bind tightly to albumin, lowering free ionized calcium and triggering positive Trousseau and Chvostek signs; your priority treatment is always isotonic sodium chloride fluid replacement to restore chloride.

MEMORY LINE

Vomiting and diuretics, tingling fingers, pH ↑ • HCO_3^- ↑

3. Retaining Ray

Respiratory Acidosis



COPD, CO₂ narcosis, pH ↓ · PaCO₂ ↑

Ray has carried a barrel chest and a pack-a-day habit for forty years. When bronchitis hits, the air goes in but refuses to come out, and the CO₂ that builds up puts him quietly to sleep.

WHY THIS CHARACTER MATTERS

Ray cannot get carbon dioxide out. Barrel chest, pursed-lip breathing and a slow slide into somnolence as CO₂ climbs. Labs: pH < 7.35 with PaCO₂ > 45 and a compensating bicarbonate. Support ventilation — sedatives and high-flow oxygen without monitoring make him worse.

The story arc

1. The setup

Forty years of smoking leave Ray with air trapped behind narrowed, floppy airways. He breathes through pursed lips to keep them open.

2. The turn

A chest infection thickens his secretions; he stops finishing sentences and his headache starts in the morning.

3. The crisis

Someone turns his oxygen to 6 L/min. An hour later he is drowsy, flushed and flapping his hands.

4. The save

Oxygen is dialed back, BiPAP goes on, secretions are cleared, and the nurse holds the ordered sedative.

3. Retaining Ray • What goes wrong

Respiratory Acidosis



What goes wrong

Acute bronchitis thickens secretions into plugs his floppy airways cannot clear; ventilation fails faster than compensation can follow. Then unregulated high-flow oxygen overshoots his saturation and suppresses the drive that was keeping him breathing.

NORMAL — THE BASELINE

Ray lives in chronic compensation: his kidneys steadily retain bicarbonate to balance his mildly acidic blood, and pursed-lip breathing keeps his airways open. He works his freight-yard shifts, sentences finished, headaches absent.

Closer look

Airway exit — Air enters Ray's lungs with little resistance, but stale gas refuses to leave: expired carbon dioxide stays trapped behind collapsed, floppy bronchioles. He exhales through pursed lips to keep the fragile passages open — and bronchitis mucus plugs narrow the exit further.

Carbon dioxide pool — The pool rises: PaCO₂ spikes to 68 millimeters of mercury while his kidneys compensate by holding bicarbonate at 32 milliequivalents per liter. His barrel chest and forty years of smoking built the trap; the infection filled it.

pH gauge — The gauge reads respiratory: pH plunged to 7.26 — below 7.35 — paired with PaCO₂ > 45 and a compensating bicarbonate. Low pH plus high PaCO₂ is always Ray's profile.

Alertness — Retained CO₂ puts him quietly to sleep: a throbbing morning headache from cerebral vasodilation, then drowsiness, flushing, and flapping asterixis hands as narcosis deepens. The cruelest trigger is kindness — someone turned his oxygen to 6 L/min, blunting his hypoxic drive.

3. Retaining Ray • Warning signs

Respiratory Acidosis



Warning signs

- pH 7.26 with PaCO₂ 68 millimeters of mercury
- Compensating bicarbonate 32 milliequivalents per liter
- PaO₂ 60 millimeters of mercury
- Morning frontal headache
- Stops finishing sentences before pausing for breath
- Tripod position, accessory muscles straining
- Drowsy, somnolent stupor — CO₂ narcosis
- Flushed and warm from peripheral vasodilation
- Bilateral flapping tremor — asterixis
- Heart rate one hundred and eighteen with frequent premature ventricular complexes
- Barrel chest and pursed-lip breathing

Priority action

Support ventilation: high-Fowler position, oxygen dialed back and titrated strictly between 88 and 92 percent saturation — never overshoot a chronic retainer's target — and urgent BiPAP to blow off retained carbon dioxide if he can protect his airway and follow commands — a patient too stuporous to protect his airway needs intubation instead, not a mask — with inhaled albuterol and ipratropium through the circuit to clear secretions. Hold every sedative and opioid — respiratory depressants make him worse — draw frequent blood gases, and intubate without delay if pH drops below 7.25 or he cannot protect his airway.

3. Retaining Ray · Check yourself

Respiratory Acidosis



Practice question

Ray has long-standing COPD. His arterial blood gas results are: pH 7.31, PaCO₂ 68 mmHg, HCO₃⁻ 32 mEq/L. The nurse interprets these results as:

- A. Metabolic acidosis without compensation
- B. Metabolic alkalosis with respiratory compensation
- C. Respiratory acidosis with partial metabolic compensation
- D. Respiratory alkalosis without compensation

ANSWER: C

The low pH with a high PaCO₂ identifies a primary respiratory acidosis — Ray retains carbon dioxide because he cannot ventilate it off. The elevated bicarbonate shows the kidneys are compensating, but the pH is still below normal, so compensation is only partial. A metabolic disorder would be driven by the bicarbonate value moving in the same direction as the pH change, which is not the primary problem here.

Remember it

To master respiratory acidosis for the NCLEX, you must fix Retaining Ray firmly in your mind. Picture his broad barrel chest, his flapping asterixis hands, and his morning headache to immediately connect hypercapnia with cerebral vasodilation and motor disturbance. Remember that Ray retains carbon dioxide, so his laboratory profile will always spotlight a low pH paired with an elevated PaCO₂, alongside a secondary compensatory rise in bicarbonate. On your exam, remember the cardinal clinical rule: never suppress a chronic retainer's ventilatory drive by overshooting their target oxygen saturation beyond ninety-two percent, and never administer respiratory depressants like benzodiazepines or opioids. When ventilation fails, your absolute priority is clearing trapped gas, supporting airway clearance, and applying noninvasive positive pressure before acidosis becomes lethal.

MEMORY LINE

COPD, CO₂ narcosis, pH ↓ · PaCO₂ ↑

4. Panicking Pam

Respiratory Alkalosis



Hyperventilation, numb lips, $pH \uparrow \cdot PaCO_2 \downarrow$

Pam's first panic attack lands her in the emergency department breathing forty times a minute. Her lips go numb, her hands claw shut, and the triage nurse has to decide whether this is fear or something in her lungs.

WHY THIS CHARACTER MATTERS

Pam blows off too much CO_2 — anxiety, pain, fever, sepsis or early PE. Numb lips, lightheadedness, carpal spasm. Labs: $pH > 7.45$, $PaCO_2 < 35$. Coach her breathing, but never assume anxiety before ruling out hypoxia and pulmonary embolism.

The story arc

1. The setup

A stressful week ends with chest tightness and the sense she cannot get enough air, so she breathes faster.

2. The turn

Her lips and fingertips go numb; she feels lightheaded and her hands curl into spasm.

3. The crisis

The nurse notices her SpO_2 is 89% on room air — a detail that does not fit a simple panic attack.

4. The save

Oxygen goes on, a D-dimer and CT angiogram are ordered, and coached slow breathing is used only after hypoxia is excluded.

4. Panicking Pam • What goes wrong

Respiratory Alkalosis



What goes wrong

A stressful week ends with chest tightness and the sense she cannot get enough air, so she breathes faster — blowing off too much CO₂. Her lips go numb, her hands claw shut, and the triage nurse has to decide whether this is fear or something in her lungs.

NORMAL — THE BASELINE

Breathing stays at a normal rate and depth, CO₂ stays in balance, and lips and hands feel normal.

Closer look

Breath rate and carbon dioxide stream — Pam blows off too much CO₂, breathing forty times a minute; rapid breaths carry carbon dioxide out in a widening stream. Causes: anxiety, pain, fever, sepsis or early PE.

pH seesaw — Labs: pH > 7.45, PaCO₂ < 35 — Panic drops the PaCO₂ and shifts the pH up into alkalosis.

Numb lips and fingertips — Her lips and fingertips go numb and she feels lightheaded — paresthesias from shifting calcium.

Clawed hands — Her hands claw shut and curl into spasm — carpal spasm from the alkalotic shift.

Oxygen saturation and embolism branch — Her SpO₂ is 89% on room air — a detail that does not fit a simple panic attack. Never assume anxiety before ruling out hypoxia and pulmonary embolism; hypoxia, sepsis, and pulmonary embolism trigger identical tachypnea.

4. Panicking Pam • Warning signs

Respiratory Alkalosis



Warning signs

- Breathing forty times a minute (hyperventilation)
- Numb lips and fingertips
- Lightheadedness
- Hands clawed shut in carpal spasm
- Chest tightness and the sense she cannot get enough air
- SpO₂ 89% on room air
- Labs: pH > 7.45, PaCO₂ < 35

Priority action

Check pulse oximetry and secure oxygenation first — oxygen goes on, and a D-dimer and CT angiogram are ordered; coached slow breathing is used only after hypoxia is excluded. Never attempt re-breathing maneuvers until underlying physiological causes of tissue hypoxia are completely ruled out.

4. Panicking Pam • Check yourself

Respiratory Alkalosis



Practice question

Pam is hyperventilating with SpO₂ 89% on room air. Which action takes priority?

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it

Remember Panicking Pam whenever you picture a patient blowing off carbon dioxide until their lips tingle and their hands claw shut. Let her name remind you that Panic drops the PaCO₂, shifts the pH up into Alkalosis, and causes Muscle spasms and paresthesias from shifting calcium. Her gasping breath mirrors the classic ABG signature of high pH and low PaCO₂. On the NCLEX, examiners love to bait you into treating hyperventilation as simple anxiety with a paper bag or a sedative. Always remember that hypoxia, sepsis, and pulmonary embolism trigger identical tachypnea. Your priority must always be checking pulse oximetry and securing oxygenation first. Never attempt re-breathing maneuvers until you have completely ruled out underlying physiological causes of tissue hypoxia.

MEMORY LINE

Hyperventilation, numb lips, pH ↑ • PaCO₂ ↓

5. Potassium Pete

Hyperkalemia



Peaked T waves, muscle cramps, hold the bananas

Pete eats bananas by the crate and his kidneys quit taking deliveries. Potassium piles up until his ECG grows tall, pointed T waves like warning spikes on a fence.

WHY THIS CHARACTER MATTERS

Pete's K^+ is above 5.0 and his heart shows it — peaked T waves, widened QRS, bradycardia. Expect muscle cramps, weakness and GI hyperactivity. Stabilize the myocardium with calcium gluconate, shift with insulin plus D50, then remove with kayexalate or dialysis.

The story arc

1. The setup

Pete has stage 4 kidney disease, takes lisinopril and spironolactone, and loves salt substitute — which is potassium chloride.

2. The turn

He notices cramping calves, tingling around his mouth, and legs that feel heavier every day.

3. The crisis

The monitor shows tall, tented T waves that slide into a widening QRS. His K^+ is 7.1.

4. The save

Calcium gluconate goes in first to armor the heart, then insulin with D50 to shift, then dialysis to actually remove it.

5. Potassium Pete • What goes wrong

Hyperkalemia



What goes wrong

Renal failure plus ACE inhibitor plus potassium-sparing diuretic plus potassium-chloride salt substitute stack into a lethal cocktail while he self-manages in the wrong direction. Metabolic acidosis pushes even more cellular potassium outward, and neuromuscular depression marches upward toward his vital organs.

NORMAL — THE BASELINE

Potassium lives primarily inside Pete's cells, and his kidneys excrete the excess from his fruit-terminal diet without drama. His ECG is unremarkable and his legs carry him through evening walks.

Closer look

Potassium field — Pete's K^+ measures a catastrophic 7.1 mEq/L, well past the critical threshold of 6.5 mEq/L and far above 5.0. The crate is literal: bananas by the crate, plus potassium chloride salt substitute dusted over every meal.

Kidney exit — His stage 4 kidneys quit taking deliveries, and two prescriptions block the exit further: lisinopril and spironolactone both stop his remaining nephrons from shedding potassium. His BUN is 68 mg/dL and creatinine 4.2 mg/dL — profound renal failure backing everything up.

Muscle cue — High potassium irritates muscles first, then paralyzes them: painful calf cramps, tingling around his mouth and fingertips, a hyperactive rumbling gut with loose diarrhea — then legs growing heavier and more rubbery until they give out entirely in ascending flaccid weakness.

ECG T wave — The monitor announces the emergency: tall, tented, peaked T waves like warning spikes on a fence, flattened P waves, and a QRS widening into a deadly slur, with heart rate down to forty-four. A widening QRS means impending cardiac arrest — reach for calcium gluconate first.

5. Potassium Pete • Warning signs

Hyperkalemia



Warning signs

- K^+ 7.1 mEq/L — above 5.0, past the 6.5 critical threshold
- Tall, peaked, tented T waves
- Flattened P waves and widening QRS
- Bradycardia of forty-four beats per minute
- Muscle cramps seizing his calves
- Tingling around the mouth and fingertips
- GI hyperactivity — cramping and loose diarrhea
- Legs heavier and more rubbery every day
- Ascending flaccid weakness — legs giving out
- BUN 68 mg/dL, creatinine 4.2 mg/dL

Priority action

Stabilize, shift, remove — in that order. Stabilize the myocardium with IV calcium gluconate first: it does not lower potassium at all; it raises the cardiac threshold and armors the heart against fibrillation. Shift with ten units of regular insulin plus fifty percent dextrose (and albuterol nebulizers), holding his lisinopril and spironolactone. Remove with sodium polystyrene sulfonate and urgent hemodialysis.

5. Potassium Pete • Check yourself

Hyperkalemia



Practice question

Pete is being discharged after treatment for hyperkalemia related to kidney failure. Which action in his discharge plan requires further teaching?

- A. Reading food labels for potassium content
- B. Avoiding potassium-rich foods such as bananas and oranges
- C. Reporting new muscle weakness or palpitations promptly
- D. Using a potassium-based salt substitute freely to season his food

ANSWER: D

Many salt substitutes are made of potassium chloride, so using one freely can quickly raise Pete's potassium again — this is a classic hidden source of potassium. Reading labels, avoiding potassium-rich foods, and reporting weakness or palpitations are all appropriate. Clients with hyperkalemia must be specifically warned about salt substitutes, not just obvious foods.

Remember it


When you remember Potassium Pete, picture him stacking crates of bananas while his ECG spikes tall, peaked T waves like sharp mountain tents. Remember that high potassium irritates muscles first with cramps and diarrhea, then paralyzes them with ascending flaccid weakness. For the NCLEX, you must always memorize the intervention sequence: stabilize the heart, shift the potassium, and remove the excess. Never forget the cardinal testing trap that intravenous calcium gluconate does not lower the serum potassium level; its sole life-saving purpose is to raise the cardiac threshold and protect the myocardium from ventricular fibrillation. When an exam question presents a patient with peaked T waves or a widening QRS complex, your hands must reach for calcium gluconate before you reach for insulin, dextrose, or gut-binding resins.

MEMORY LINE

Peaked T waves, muscle cramps, hold the bananas

6. Low-K Lena

Hypokalemia



Flat T waves, U waves, silent belly, never IV push

Lena takes her water pill faithfully and eats almost nothing. Her heart's tracing flattens out, her bowel goes silent, and a U wave shows up on the monitor like a signature at the end of every beat.

WHY THIS CHARACTER MATTERS

Lena's K⁺ is below 3.5 from diuretics, vomiting or NG suction. Everything gets flat and quiet: flat T waves, U waves, weak muscles, absent bowel sounds. Replace potassium slowly — diluted, on a pump, never IV push, never faster than 10 mEq/hr on a general floor.

The story arc

1. The setup

Furosemide daily for heart failure, a week of poor appetite, and no potassium supplement refilled.

2. The turn

Her legs feel rubbery on the stairs, her belly bloats, and the nurse cannot hear bowel sounds.

3. The crisis

The monitor shows flattened T waves with a U wave, then a run of PVCs. She also takes digoxin.

4. The save

Diluted KCl runs on a pump through a good IV site with continuous ECG monitoring — never a push, never undiluted.

6. Low-K Lena • What goes wrong

Hypokalemia



What goes wrong

A viral stomach bug kills her appetite for three weeks while the loop diuretic keeps draining potassium she never replaces. Excitable tissues hyperpolarize and go sluggish system-wide until she can barely lift her arms and her breathing grows shallow.

NORMAL — THE BASELINE

With potassium replaced alongside her diuretic and regular meals of potatoes, bananas, and spinach, Lena's membrane potentials hold steady. Her tracing, bowel, and legs all keep their normal tone.

Closer look

Potassium field — Lena's K^+ is below 3.5 — a severe 2.4 mEq/L — drained by daily furosemide for heart failure, three weeks of eating almost nothing, and a potassium supplement she never refilled. The organizer's empty slots are the story: the water pill was taken faithfully, the potassium was not.

T/U waves — Low potassium makes everything low, slow, and flat on the tracing: flattened T waves, distinct ST depression, and a prominent U wave trailing each QRS like a signature at the end of every beat. Then come frequent premature ventricular contractions.

Bowel motion — Her bowel goes silent: a completely silent abdomen on auscultation means paralytic ileus, with bloating, hypoactive reflexes, and rubbery legs that buckle on the stairs. Oral supplements wait until bowel sounds safely return.

Medication source — Two medication traps hide here: her daily digoxin becomes dangerous because low potassium leaves digitalis receptors uninhibited, heightening lethal ectopy risk. And refractory hypokalemia never corrects until magnesium is replaced first — her magnesium is 1.2 mg/dL.

6. Low-K Lena • Warning signs

Hypokalemia



Warning signs

- K^+ 2.4 mEq/L — below the 3.5 floor
- Flat T waves with ST depression
- Prominent U waves after every beat
- Frequent premature ventricular contractions
- Silent belly — absent bowel sounds, paralytic ileus
- Abdominal bloating
- Rubbery, heavy legs and nocturnal calf cramps
- Hypoactive deep tendon reflexes
- Barely able to lift her arms; shallow breathing
- Chronic exhaustion, unquenchable thirst, polyuria
- Magnesium 1.2 mg/dL — and she takes digoxin

Priority action

Replace potassium slowly — diluted IV potassium chloride on an infusion pump at exactly 10 mEq per hour peripherally, never faster on a general floor, with continuous ECG monitoring and vigilance for chemical phlebitis at the site. Never IV push, never undiluted: an unfiltered surge stops the heart instantly. Replace magnesium concurrently, and plan oral supplements with meals once bowel sounds return.

6. Low-K Lena • Check yourself

Hypokalemia



Practice question

Lena's potassium remains 2.9 mEq/L despite several rounds of IV potassium replacement. Her magnesium level returns at 1.1 mg/dL. Which action should the nurse anticipate next?

- A. Replace her magnesium, because hypokalemia will not correct until the magnesium deficit is treated
- B. Increase the IV potassium rate and give it by rapid infusion
- C. Stop all potassium replacement and recheck the level in one week
- D. Restrict her dietary potassium until the serum level stabilizes

ANSWER: A

Hypokalemia that resists replacement is commonly driven by an uncorrected magnesium deficit; the potassium will not hold until magnesium is replaced. Speeding up IV potassium is dangerous — potassium is never given IV push and is infused slowly on a pump. Dietary potassium restriction is the opposite of what a hypokalemic client needs.

Remember it

To lock in Low-K Lena on your exams, remember that low potassium makes everything low, slow, and flat. Picture Lena's heart tracing flattening out into collapsed T waves while a looping U wave signs her name at the end of each beat. Envision her silent belly completely paralyzed by ileus and her rubbery legs buckling under weak reflexes. Above all, stamp the absolute rule into your memory: never administer potassium by intravenous push or bolus, because an unfiltered surge will stop a patient's heart instantly. For the NCLEX, always check the patient's magnesium level when hypokalemia resists standard replacement therapy; refractory potassium deficits will never correct until you replace magnesium first. Keep Lena's pump slow and steady, never exceeding ten milliequivalents hourly peripherally.

MEMORY LINE

Flat T waves, U waves, silent belly, never IV push

7. Salty Sal

Hypernatremia



Thirsty, dry, agitated — $\text{Na}^+ > 153$, correct it slowly

Sal is a nursing-home resident on tube feeds who can no longer say he is thirsty. Salt quietly concentrates in his blood until his tongue is like sandpaper and he is picking at the sheets.

WHY THIS CHARACTER MATTERS

Sal has more sodium than water — dehydration, tube feeds without free water, or diabetes insipidus. Thirst, dry sticky mucous membranes, restlessness and twitching. Correct slowly with hypotonic fluid: too fast and the brain cells swell.

The story arc

1. The setup

Continuous tube feeds run for days with no free-water flushes, and Sal cannot verbalize thirst.

2. The turn

His tongue is furrowed and sticky, his urine is scant and dark, and he grows restless and irritable.

3. The crisis


Na^+ returns at 162. He twitches, then has a brief seizure.

4. The save

Hypotonic fluid is titrated to drop sodium no faster than about 0.5 mEq/L per hour, with free-water flushes added to the feeding plan.

7. Salty Sal • What goes wrong

Hypernatremia



What goes wrong

His formula rate increases for calories while the free-water flushes are inadvertently omitted for days. The deficit expands silently because he cannot voice his thirst, and cellular dehydration steals moisture from his brain until delirium and sheet-picking announce the crisis.

NORMAL — THE BASELINE

With scheduled free-water flushes matching his tube feeds, Sal's sodium stays within its normal range of 135 to 145 mEq/L and his neurons stay hydrated. The deli master who mixed salt brines by hand is cared for by a plan that respects the salt-to-water balance.

Closer look

Sodium concentration — Sal has more sodium than water: his serum sodium climbs above 145 to a dangerous 162 mEq/L, with serum osmolality surged to 328 mOsm/kg. The feeding pump concentrates it further — hyperosmolar formula without free water draws fluid from his tissues to process the solute load.

Neuron water — Hypertonic blood pulls water out of his brain cells until they shrink under osmotic tension: irritability, fidgeting, and sleep fragmentation first, then twitching and a brief generalized tonic-clonic seizure as shrunken neurons fire abnormally. Correct too fast and the same cells swell — cerebral edema runs both directions.

Mouth/thirst cues — Thirsty, dry, agitated — but Sal cannot say he is thirsty: post-stroke expressive aphasia silences the alarm. His tongue becomes a rough, sandpaper-dry slab with deep longitudinal furrows, his lips crack, and his kidneys conserve scant, dark, concentrated urine.

Correction speed — The guardrail is numeric: drop sodium no faster than about 0.5 mEq/L per hour, and no more than ten to twelve mEq/L in twenty-four hours. Neuro checks every hour, intake and output tracked, repeat sodium every four hours.

7. Salty Sal · Warning signs

Hypernatremia



Warning signs

- Na^+ 162 — climbing above 145
- Serum osmolality 328 mOsm/kg
- Tongue furrowed, sticky, and sandpaper-dry
- Dry, cracked lips and parched oral mucosa
- Scant, dark, concentrated urine
- Restlessness, irritability, and fidgeting
- Picking at the bed sheets, acute confusion and delirium
- Skin tenting with severe poor turgor
- Coarse muscle twitches across the forearms
- Brief generalized tonic-clonic seizure
- Elevated BUN-to-creatinine ratio of thirty to one

Priority action

Correct slowly with hypotonic fluid: 0.45% sodium chloride titrated to drop sodium no faster than about 0.5 mEq/L per hour, with scheduled free-water flushes added back to the feeding plan. Never slam in free water — rapid shifts flood dried-out neurons and cause catastrophic cerebral edema. Seizure precautions with padded rails, lateral positioning, and suction protect him while the numbers come down.

7. Salty Sal • Check yourself

Hypernatremia



Practice question

Sal, a nonverbal nursing-home resident receiving tube feedings, was treated for hypernatremia with a sodium of 162 mEq/L. Which care-plan intervention best prevents recurrence?

- A. Restrict all water flushes to keep his sodium from falling
- B. Provide scheduled free-water flushes as ordered and monitor intake, output, and daily weights
- C. Add salt to his tube feedings so his serum sodium stays stable
- D. Check his sodium only when he becomes restless or agitated

ANSWER: B

Tube-fed clients who cannot communicate thirst depend on staff for free water; scheduled free-water flushes with intake, output, and weight monitoring prevent the water deficit that causes hypernatremia. Restricting flushes would recreate the problem, and adding salt would worsen it. Waiting for restlessness means waiting for a symptom of brain-cell dehydration rather than preventing it.

Remember it

Remember Salty Sal as the tube-fed resident whose cracked, sandpaper tongue and agitated sheet-picking warn you of a deadly salt-to-water mismatch. Picture his name to immediately recall hypernatremia: a serum sodium climbing above 145 mEq/L caused by unreplaced free water in patients who cannot voice intense thirst. Always associate Sal's twitching muscles and sudden seizures with shrinking, dehydrated brain cells begging for hydration. On your NCLEX exam, remember this golden rule of fluid management: correct hypernatremia slowly and deliberately using hypotonic fluids. Never slam intravenous free water or drop serum sodium too rapidly, because shifting fluid into dried-out neurons causes catastrophic cerebral edema and brain herniation. When caring for nonverbal tube-fed clients, your diligent free-water flushes will keep your patients from becoming Salty Sal.

MEMORY LINE

Thirsty, dry, agitated — $\text{Na}^+ > 153$, correct it slowly

8. Diluted Dana

Hyponatremia



Too much water, $\text{Na}^+ < 135$, headache to seizure

Dana's body will not let go of a single drop. Her urine is dark and scarce while her blood grows more dilute by the hour, and the first thing that swells is her brain.

WHY THIS CHARACTER MATTERS

Dana holds onto water she does not need — SIADH from small-cell lung cancer, head injury or SSRIs. Sodium falls, brain cells swell: headache, nausea, confusion, seizures. Fluid restriction is the core treatment; 3% saline only for severe symptomatic lows, and only slowly.

The story arc

1. The setup

Dana is treated for small-cell lung cancer and started on an SSRI. Her weight climbs two kilograms with no edema.

2. The turn

She develops a dull headache, nausea and a sense that her thoughts are slow.

3. The crisis

Sodium is 116 and she has a generalized seizure in the hallway.

4. The save

3% saline runs on a pump with hourly sodium checks and strict fluid restriction to 800 mL/day.

8. Diluted Dana • What goes wrong

Hyponatremia



What goes wrong

Small-cell lung cancer secretes ectopic antidiuretic hormone, and a newly started SSRI amplifies the release further. Over several days she dilutes silently — two kilograms gained without a trace of edema — while her brain cells imbibe water and begin to swell.

NORMAL — THE BASELINE

Dana's watercolor washes stay on the paper: antidiuretic hormone rises and falls appropriately, her sodium holds at or above 135 mEq/L, and her sharp illustrator's mind stays clear. Weight, urine, and thirst all balance.

Closer look

Water volume — Dana's body will not let go of a single drop: continuous antidiuretic signals command her collecting ducts to hoard pure water. Her weight climbs two kilograms — with no edema, no crackles, and no jugular distention, because the water spreads evenly through every compartment.

Sodium particles — Too much water and her sodium dilutes below 135: a critical 116 mEq/L, beneath the danger threshold of 120, with serum osmolality profoundly depressed at 242 mOsm/kg. Headache to seizure is the trajectory as the number falls.

Urine concentration — The specimen cup holds the paradox: while her blood dilutes, her kidneys produce dark, scarce, concentrated urine — urine osmolality of 680 mOsm/kg with urine sodium of 52 mEq/L. The kidneys are obeying the hormone, not the blood.

Brain — The first thing that swells is her brain: hypoosmolar water shifts into her cerebral cortex, producing the downward spiral from dull frontal headache and nausea to slow thoughts, disorientation, and a generalized seizure in the hallway. Cerebral edema is what makes dilution deadly.

8. Diluted Dana • Warning signs

Hyponatremia



Warning signs

- Sodium 116 mEq/L — below 135, beneath the 120 danger threshold
- Serum osmolality 242 mOsm/kg
- Concentrated urine — osmolality 680 mOsm/kg, urine sodium 52 mEq/L
- Weight gain of two kilograms with no edema
- Dull frontal headache behind the eyes
- Persistent nausea and generalized malaise
- Thoughts slow — sluggish indifference
- Profound muscle weakness and acute disorientation
- Generalized tonic-clonic seizure, then unresponsive coma

Priority action

Fluid restriction is the core treatment — strict restriction to 800 mL/day, hypotonic fluids discontinued, strict intake and output with daily morning weights. For severe symptomatic lows like Dana's seizure, run 3% saline slowly on a dedicated pump with serial sodium checks — and never correct faster than 8 to 12 mEq/L in twenty-four hours, or osmotic demyelination can lock her into central pontine paralysis. Seizure precautions with padded rails and suction complete the picture.

8. Diluted Dana • Check yourself

Hyponatremia



Practice question

Dana with SIADH is receiving hypertonic saline for severe symptomatic hyponatremia. Her sodium has risen from 112 mEq/L to 125 mEq/L in 10 hours. What should the nurse do?

- A. Increase the infusion rate so the sodium normalizes by the end of the shift
- B. Stop the infusion and notify the provider, because the sodium is correcting too rapidly
- C. Encourage free water by mouth to slow the rise in sodium
- D. Document the result as an expected, routine rate of correction

ANSWER: B

A rise of 13 mEq/L in 10 hours exceeds the safe correction limit of about 8 to 12 mEq/L in 24 hours and risks osmotic demyelination syndrome, so the infusion should be stopped and the provider notified. Normalizing sodium within a shift is exactly the error this question tests. Oral free water is also inappropriate in SIADH, where fluid restriction is a core treatment.

Remember it

Remember Diluted Dana whenever you face fluid balance questions on the NCLEX. Picture Dana drowning in her own clear watercolor wash: her body holds too much pure water, causing dilutional hyponatremia with serum sodium plummeting below 135 mEq/L while her weight rises without peripheral edema. Her swollen brain creates the classic downward spiral progressing from a dull headache and nausea to disorientation and fatal seizures. For your exam, always remember this cardinal rule: in SIADH, the primary intervention is strict fluid restriction, but when severe symptoms strike, hypertonic 3% saline must be infused slowly and cautiously on a dedicated pump. Never correct sodium faster than 8 to 12 mEq/L in a twenty-four-hour window, or you risk devastating osmotic demyelination syndrome.

MEMORY LINE

Too much water, $Na^+ < 135$, headache to seizure

9. Chvostek Chuck

Hypocalcemia



Twitchy face, tingling lips, watch the airway

Chuck wakes from thyroid surgery with a buzzing lip and a cheek that jumps when the nurse taps it. His parathyroids took the hit, and now every nerve in his body has its volume turned up.

WHY THIS CHARACTER MATTERS

Chuck's calcium falls after thyroid surgery, from chronic kidney disease or from massive transfusion. Everything gets irritable: facial twitching (Chvostek's), carpal spasm (Trousseau's), tingling and, at worst, laryngospasm. Keep calcium gluconate and a trach tray nearby.

The story arc

1. The setup

A total thyroidectomy leaves his parathyroid glands bruised and underperforming.

2. The turn

He reports tingling around his mouth and in his fingertips overnight.

3. The crisis

His cheek twitches when the nurse taps his facial nerve, his hand claws under the cuff, and then his voice turns high and crowing.

4. The save

IV calcium gluconate is pushed slowly on a monitor, with a tracheostomy tray at the bedside.

9. Chvostek Chuck • What goes wrong

Hypocalcemia



What goes wrong

Thirty-six hours after total thyroidectomy, his bruised parathyroids underperform and the circulating calcium pool drains. Every nerve's volume turns up: paresthesias first, then tetany, then the airway itself joins the spasm.

NORMAL — THE BASELINE

With calcium in range, Chuck's nerves stay properly insulated and calm, firing only when told. The drama coach talks with his hands and twitches his cheek on purpose, as a cue to the backstage crew.

Closer look

Calcium gauge — Chuck's total serum calcium is 6.8 mg/dL, well beneath the normal floor of 8.5 mg/dL, with a critically low ionized fraction. His thyroidectomy bruised and devascularized his parathyroid glands, cutting off the hormone that resorbs bone, reclaims calcium in the kidney, and activates vitamin D. Chronic kidney disease or massive citrated transfusion can drop calcium the same way.


Facial nerve — Tap the facial nerve just anterior to the ear and his whole cheek winces into spasm: a positive Chvostek's sign, the drama coach's stage trick turned diagnostic. Low calcium destabilizes neuronal membranes so sodium rushes in and every threshold falls.

Lips/hands — The letter C carries his signs: circumoral tingling — lips buzzing like tiny bees — and carpal spasm, the hand clawing into involuntary flexion when the blood pressure cuff inflates, a positive Trousseau's sign. Pins-and-needles creep through his fingertips and deep cramps seize his calves, with wildly brisk patellar reflexes.

Airway watch — The crowing gasp is the emergency: laryngospasm constricts his vocal cords into high-pitched stridor. Keep a tracheostomy tray within arm's reach post-thyroidectomy, keep calcium gluconate ready as the rapid antidote, and never leave him unattended while he stridors.

9. Chvostek Chuck • Warning signs

Hypocalcemia



Warning signs

- Tingling around the mouth — buzzing lips
- Fingertip paresthesias
- Cheek twitch when the facial nerve is tapped — Chvostek's sign
- Hand claws under the blood pressure cuff — Trousseau's sign
- Hyperactive deep tendon reflexes
- Deep muscle cramps in the calves
- High, crowing voice — laryngospasm and stridor
- Total calcium 6.8 mg/dL — beneath the 8.5 floor
- Prolonged QT interval on the twelve-lead ECG

Priority action

Protect the airway first: call for help, position him upright, and bring the tracheostomy tray and suction to the bedside. Then give IV calcium gluconate slowly through a dedicated line under continuous cardiac monitoring — verify telemetry before infusing, because pushing it too quickly causes bradycardia, hypotension, and cardiac standstill. Add seizure precautions, padded rails, dimmed lights, magnesium checks, and watch the QT interval narrow as he recovers.

9. Chvostek Chuck · Check yourself

Hypocalcemia



Practice question

One day after thyroid surgery, Chuck reports circumoral tingling. While the nurse inflates a blood pressure cuff on his arm and leaves it inflated, Chuck's hand and fingers go into a spasm. The nurse interprets this Trousseau's sign as indicating:

- A. Hyperkalemia
- B. Hypercalcemia
- C. Hyponatremia
- D. Hypocalcemia

ANSWER: D

Carpal spasm with cuff inflation (Trousseau's sign), like facial twitching with tapping (Chvostek's sign), reflects the neuromuscular irritability of hypocalcemia, a risk after thyroid surgery if the parathyroids are injured. Hypercalcemia produces the opposite picture of decreased neuromuscular excitability. Calcium gluconate and emergency airway equipment should be kept nearby because severe hypocalcemia can cause laryngospasm.

Remember it

You can always remember hypocalcemia by picturing Chvostek Chuck clutching his buzzing lips while his cheek jumps to the nurse's tap. Connect the letter C in Chuck to Chvostek's Cheek twitch, Circumoral tingling, Carpal spasm under the blood pressure cuff, and Calcium gluconate as the rapid antidote. Whenever you think of Chuck gasping his theater lines, remember that his crowing stridor means laryngospasm, requiring a tracheostomy tray within arm's reach post-thyroidectomy. For your NCLEX exam, always remember that an acute drop in calcium produces neuromuscular irritability, hyperactive deep tendon reflexes, and a prolonged QT interval. When answering priority questions, never leave a hypocalcemic client unattended if they develop stridor, and always verify continuous telemetry before infusing IV calcium gluconate because pushing it too quickly causes fatal cardiac arrest.

MEMORY LINE

Twitchy face, tingling lips, watch the airway

10. Stony Stella

Hypercalcemia



Stones, bones, groans and psychiatric overtones

Stella turns to stone from the inside out. Calcium leaches out of her bones, settles in her kidneys and slows every muscle and thought she has.

WHY THIS CHARACTER MATTERS

Stella's calcium is above 10.5 from malignancy or hyperparathyroidism. Everything slows: constipation, lethargy, weak reflexes, kidney stones and bone pain. Isotonic fluids first, then calcitonin and bisphosphonates; keep her moving and safe from fractures.

The story arc

1. The setup

Metastatic breast cancer erodes bone and dumps calcium into her blood while she stays in bed for weeks.

2. The turn

She becomes constipated, drinks constantly, urinates constantly, and her family says she is 'not herself.'

3. The crisis

Calcium is 14.2, she is lethargic and her ECG shows a shortened QT with bradycardia.

4. The save

Isotonic saline restores volume and promotes excretion, calcitonin works fast, and a bisphosphonate holds the line.

10. Stony Stella • What goes wrong

Hypercalcemia



What goes wrong

Metastatic breast cancer erodes bone and dumps calcium into her blood while she stays in bed for weeks. She becomes constipated, drinks constantly, urinates constantly, and turns to stone from the inside out.

NORMAL — THE BASELINE

When calcium stays in range — at or below 10.5 mg/dL — it does not leach from bone, settle in kidneys, or slow every muscle and thought.

Closer look

Kidney stones — Calcium leaches out of her bones and settles in her kidneys — stones are the first word of the rhyme: stones, bones, groans and psychiatric overtones.

Bone pain — Bones that ache and break as calcium leaves them; keep her moving and safe from fractures.

Groans and constipation — Everything slows: constipation, lethargy, weak reflexes — groans from constipation and nausea.

Psychiatric slowing — Psychiatric overtones from lethargy and confusion; her family says she is 'not herself.'

Shortened QT and saline first — Her ECG shows a shortened QT with bradycardia. In an acute hypercalcemic crisis, the first priority is aggressive intravenous hydration with 0.9% normal saline to flush the mineral out through the renal tubules. Never pick a thiazide diuretic — thiazides reabsorb calcium rather than excrete it.

10. Stony Stella • Warning signs

Hypercalcemia



Warning signs

- Kidney stones
- Bone pain
- Constipation
- Nausea
- Lethargy
- Confusion ('not herself')
- Weak (depressed) reflexes
- Constant thirst and constant urination
- Shortened QT interval
- Bradycardia
- Calcium 14.2 (her crisis value)

Priority action

Isotonic fluids first — aggressive 0.9% normal saline restores volume and promotes excretion — then calcitonin, which works fast, and a bisphosphonate to hold the line. Keep her moving and safe from fractures.

10. Stony Stella • Check yourself

Hypercalcemia



Practice question

Which initial therapy does the nurse anticipate for Stella's calcium of 14.2 mg/dL?

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it

To master hypercalcemia on your clinical exams, remember Stony Stella and her famous rhyme: stones in the kidneys, bones that ache and break, groans from constipation and nausea, and psychiatric overtones from lethargy and confusion. Picture Stella as a petrified statue where high calcium slows everything down, causing bradycardia, depressed reflexes, and a shortened QT interval on the electrocardiogram. When an NCLEX question presents an acute hypercalcemic crisis, your first priority is always aggressive intravenous hydration with 0.9% normal saline to flush the mineral out through the renal tubules. Never pick a thiazide diuretic on your exam, because thiazides reabsorb calcium rather than excrete it, which will only turn a patient like Stella further into solid stone.

MEMORY LINE

Stones, bones, groans and psychiatric overtones

11. Magnesium Maggie

Magnesium Imbalance



Low Mg = torsades; high Mg = reflexes gone

Maggie is the electrolyte everyone forgets until the monitor twists into torsades. On one side of her story the reflexes are hyperactive; on the other they vanish completely.

WHY THIS CHARACTER MATTERS

Maggie swings both ways. Low magnesium (alcohol use, malnutrition, diuretics) causes tremors, hyperreflexia and torsades de pointes. High magnesium (renal failure, eclampsia infusions) wipes out reflexes and respirations — calcium gluconate is the antidote.

The story arc

1. The setup

Years of heavy alcohol use and poor intake quietly drain Maggie's magnesium stores.

2. The turn

Her hands tremor, her reflexes are brisk, and her potassium refuses to come up despite repeated replacement.

3. The crisis

The monitor shows a QT that keeps stretching, then a twisting polymorphic rhythm.

4. The save

Magnesium sulfate is infused — and the nurse then watches reflexes, respirations and blood pressure for the opposite problem.

11. Magnesium Maggie • What goes wrong

Magnesium Imbalance



What goes wrong

Years of heavy alcohol use and poor intake quietly drain Maggie's magnesium stores. Her hands tremor, her reflexes are brisk, and her potassium refuses to come up despite repeated replacement — then her QT keeps stretching into torsades.

NORMAL — THE BASELINE

Maggie swings wildly between two dangerous extremes — twitching low and sedated high. In balance, reflexes and respirations stay present and steady, and potassium replacement works.

Closer look

Low-magnesium tremor — Low magnesium (alcohol use, malnutrition, diuretics) causes tremors and hyperreflexia.

Resistant potassium — Hypokalemia that resists aggressive replacement is almost always caused by hidden hypomagnesemia — replace magnesium first, because potassium refuses to come up despite repeated replacement until magnesium is fixed.

Twisting torsades — Low magnesium provokes torsades de pointes — the monitor twists into a twisting polymorphic rhythm after a QT that keeps stretching.

High-magnesium reflex loss — High magnesium (renal failure, eclampsia infusions) wipes out reflexes and respirations. The sudden loss of a patellar reflex is the earliest warning sign of impending toxicity before respiratory arrest.

Calcium gluconate antidote — Calcium gluconate is the antidote for high magnesium. On any therapeutic magnesium infusion, assess deep tendon reflexes before touching the rate.

11. Magnesium Maggie • Warning signs

Magnesium Imbalance



Warning signs

- Tremors
- Hyperreflexia (brisk reflexes)
- QT that keeps stretching
- Torsades de pointes (twisting polymorphic rhythm)
- Loss of the patellar reflex (high-magnesium toxicity)
- Fading respirations (high magnesium)

Priority action

Infuse magnesium sulfate for the low side — and then watch reflexes, respirations and blood pressure for the opposite problem. For high magnesium, calcium gluconate is the antidote.

11. Magnesium Maggie • Check yourself

Magnesium Imbalance



Practice question

During Maggie's magnesium sulfate infusion, her patellar reflex is absent and respirations are 10/min. The nurse should:

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it


You can easily remember Magnesium Maggie by picturing her swinging wildly between two dangerous extremes: twitching low and sedated high. Link low magnesium to Maggie's spastic gymnastics, remembering that deficient magnesium provokes tremors, hyperreflexia, and that telltale twisting torsades de pointes on the cardiac strip. If Maggie receives too much magnesium, she turns into a limp ragdoll whose reflexes and respirations completely fade away. For the NCLEX exam, remember that hypokalemia that resists aggressive replacement is almost always caused by hidden hypomagnesemia, so replace magnesium first. Furthermore, whenever you manage a therapeutic magnesium infusion, always assess deep tendon reflexes before touching the rate; the sudden loss of a patellar reflex is your earliest warning sign of impending toxicity before respiratory arrest occurs.

MEMORY LINE

Low Mg = torsades; high Mg = reflexes gone

12. Septic Sam

Distributive Shock



Warm flushed skin early, then BP bottoms out

Sam throws a party in every blood vessel he owns — everything dilates, everything opens up. For an hour he's warm and pink and looks fine. Then the party drains the tank and his pressure falls through the floor.

WHY THIS CHARACTER MATTERS

Sam looks deceptively good early: warm, flushed skin and bounding pulses from massive vasodilation. Then MAP crashes, lactate climbs and urine output drops. Cultures before antibiotics, aggressive fluids, then vasopressors if MAP stays under 65.

The story arc

1. The setup

A urinary tract infection in an 78-year-old goes untreated for a week; the only complaint is that he seems confused.

2. The turn

He arrives warm, pink and bounding — heart rate 118, temperature 38.9°C, blood pressure still 'okay.'

3. The crisis


Lactate is 4.6, MAP falls to 58 after two liters of fluid, and urine output stops.

4. The save

Cultures were drawn before the first antibiotic, 30 mL/kg of crystalloid went in, and norepinephrine is now titrated to a MAP of 65.

12. Septic Sam • What goes wrong

Distributive Shock



What goes wrong

A urinary tract infection in a 78-year-old goes untreated for a week; the only complaint is that he seems confused. He arrives warm, pink and bounding with blood pressure still "okay," then the party drains the tank — MAP crashes, lactate climbs and urine output drops; a lactate ≥ 4 mmol/L with persistent hypotension despite fluid resuscitation defines septic shock.

NORMAL — THE BASELINE

Vessels hold normal tone, MAP stays adequate, lactate stays low, and urine flows.

Closer look

Vessel radius — Sam throws a party in every blood vessel he owns — everything dilates, everything opens up: massive vasodilation makes him look deceptively good early.

Resistance — As the vessels stay wide open, resistance falls and the vascular tank drains; MAP falls to 58 after two liters of fluid.

Blood pressure — Cultures before antibiotics, aggressive fluids, then vasopressors if MAP stays under 65: norepinephrine is titrated to a MAP of 65.

Skin/perfusion — Early he is warm, flushed and pink with bounding pulses — heart rate 118, temperature 38.9°C — then crashing into cold, mottled collapse once the vascular tank runs dry; lactate climbs to 4.6 and urine output stops.

12. Septic Sam • Warning signs

Distributive Shock



Warning signs

- Confusion as the only early complaint
- Warm, flushed skin and bounding pulses
- Heart rate 118, temperature 38.9°C
- Lactate 4.6
- MAP falls to 58 after two liters of fluid
- Urine output stops

Priority action

Obtain blood cultures first — drawing cultures before the antibiotic dose is non-negotiable — hang broad-spectrum IV antibiotics within one hour, push 30 mL/kg of crystalloid fluids, and start norepinephrine if MAP remains under 65.

12. Septic Sam • Check yourself

Distributive Shock



Practice question

A client with septic shock has received blood cultures, broad-spectrum IV antibiotics within the hour, and 30 mL/kg of crystalloid fluids. The MAP remains 58 mmHg and lactate is elevated. What should the nurse anticipate NEXT?

- A.** Start a vasopressor such as norepinephrine to restore MAP to at least 65 mmHg
- B.** Stop all IV fluids and reassess in 4 hours
- C.** Repeat blood cultures and wait for results before any further treatment
- D.** Administer an antipyretic and recheck the blood pressure in the morning

ANSWER: A

Persistent hypotension with MAP below 65 mmHg despite adequate fluid resuscitation defines septic shock requiring vasopressor support, with norepinephrine as the first-line agent. Waiting, stopping fluids, or delaying for culture results allows tissue hypoperfusion and rising lactate to progress.

Remember it

Remember Septic Sam as the party host whose open doors represent wide-open, dilated vessels: warm, pink, and bounding early, but crashing into cold, mottled collapse once the vascular tank runs dry. When you picture Sam, visualize his three-step lifeline to anchor the sepsis resuscitation protocol directly in your memory. You must obtain blood cultures first, hang broad-spectrum intravenous antibiotics within one hour, and push thirty milliliters per kilogram of crystalloid fluids before starting norepinephrine if the mean arterial pressure remains stubborn. On the NCLEX, examiners love testing this sequence and the diagnostic threshold for septic shock. Always remember that a lactate value greater than or equal to 4 millimoles per liter with persistent hypotension despite fluid resuscitation defines septic shock, and drawing cultures before the antibiotic dose is non-negotiable.

MEMORY LINE

Warm flushed skin early, then BP bottoms out

13. Pumpless Paul

Cardiogenic Shock



Pump fails: cold, clammy, crackles, high filling pressures

Paul's pump loses half its muscle in a single morning. The tank is full but nothing circulates — his lungs fill from behind while his fingertips go grey.

WHY THIS CHARACTER MATTERS

Paul's heart cannot move the blood it already has. After a large anterior MI he is cold, clammy and hypotensive with crackles and jugular vein distention. Fluids drown him; he needs inotropes, afterload reduction and reperfusion.

The story arc

1. The setup

An anterior STEMI is treated late; a large section of left ventricle stops contracting.

2. The turn

His skin turns cool and mottled, his pulses thread out, and he cannot lie flat without gasping.

3. The crisis

Blood pressure 78/50, crackles to the scapulae, jugular veins distended, urine output 10 mL/hr.

4. The save

Dobutamine improves contractility, careful diuresis unloads the lungs, and he goes to the cath lab for revascularization.

13. Pumpless Paul • What goes wrong

Cardiogenic Shock



What goes wrong

An anterior STEMI is treated late; a large section of left ventricle stops contracting. His lungs fill from behind while his fingertips go grey — Paul's pump loses half its muscle in a single morning.

NORMAL — THE BASELINE

A healthy ventricle moves the blood it receives forward — lungs stay clear, skin stays warm, and urine output keeps flowing.

Closer look

Weak left ventricle — After a large anterior MI, a large section of left ventricle stops contracting. Paul's heart cannot move the blood it already has — the tank is full but nothing circulates.

Wedge pressure and crackles — Picture his high wedge pressure and critically low cardiac output: crackles to the scapulae and jugular vein distention as filling pressure backs into the lungs.

Cool mottled skin — Cold, clammy and hypotensive: his skin turns cool and mottled, his pulses thread out, and he cannot lie flat without gasping while his fingertips go grey.

No large-bolus warning — Never give a large fluid bolus to a patient in cardiogenic shock. Fluids drown him — fluid resuscitating Pumpless Paul will rapidly drown him in pink, frothy sputum.

Inotrope and reperfusion path — Dobutamine improves contractility, careful diuresis unloads the lungs, and he goes to the cath lab for revascularization: prioritize positioning, oxygenation, intravenous inotropes to strengthen the pump, and rapid preparation for coronary reperfusion.

13. Pumpless Paul • Warning signs

Cardiogenic Shock



Warning signs

- Blood pressure 78/50
- Crackles to the scapulae
- Jugular vein distention
- Urine output 10 mL/hr
- Cool, mottled skin
- Thready pulses
- Cannot lie flat without gasping
- Cold and clammy

Priority action

Inotropes to strengthen the pump (dobutamine), positioning and oxygenation, careful diuresis to unload the lungs, and rapid preparation for the cath lab and reperfusion — never a large fluid bolus, and no afterload-reducing vasodilators while his pressure is this low (systolic 78).

13. Pumpless Paul • Check yourself

Cardiogenic Shock



Practice question

Paul is hypotensive with crackles and JVD after an anterior MI. Which order should the nurse question?

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it

Remember Pumpless Paul whenever you encounter heart failure transforming into cardiovascular shock: his broken pump keeps the tank full, creating cold, clammy skin, crackles to the scapulae, and elevated filling pressures. When you see Paul shivering in his waterlogged overalls with swollen neck veins, picture his high wedge pressure and critically low cardiac output. For the NCLEX exam, remember the vital distinction between cardiogenic shock and distributive shock: never give a large fluid bolus to a patient in cardiogenic shock. Fluid resuscitating Pumpless Paul will rapidly drown him in pink, frothy sputum. Instead, prioritize positioning, oxygenation, intravenous inotropes to strengthen the pump, and rapid preparation for coronary reperfusion or mechanical circulatory assist devices.

MEMORY LINE

Pump fails: cold, clammy, crackles, high filling pressures

14. Bleeding Bea

Hypovolemic Shock



Tank is empty: tachycardia first, hypotension late

Bea bleeds into her own abdomen where no one can see it. Her heart races to make up the difference for an hour, and by the time her blood pressure falls she has already lost a third of her volume.

WHY THIS CHARACTER MATTERS

Bea loses volume faster than her body can compensate — trauma, GI bleed, burns. Tachycardia and narrowing pulse pressure come first; hypotension is a late sign. Two large-bore IVs, isotonic crystalloid, blood products, and control the source.

The story arc

1. The setup

A car crash leaves her with a splenic injury and no external bleeding to point at.

2. The turn

Heart rate 124, respirations 26, skin cool and pale, restless — but blood pressure reads 112/88.

3. The crisis

Blood pressure drops to 80/50, urine output stops, and she becomes confused.

4. The save

Two large-bore IVs, warmed isotonic crystalloid, type-specific blood and surgery for source control.

14. Bleeding Bea • What goes wrong

Hypovolemic Shock



What goes wrong

Bea loses volume faster than her body can compensate — trauma, GI bleed, burns. Her heart races furiously long before the engine finally stalls, and by the time her blood pressure falls she has already lost a third of her volume.

NORMAL — THE BASELINE

With her circulating volume intact, the tank stays full — pressure, perfusion and urine output all hold.

Closer look

Occult abdominal bleeding — A car crash leaves her with a splenic injury and no external bleeding to point at. Bea bleeds into her own abdomen where no one can see it.

Tachycardia and narrowing pulse pressure — Tachycardia and narrowing pulse pressure come first — heart rate 124, respirations 26, blood pressure still reading 112/88 — the earliest reliable heralds of hypovolemic collapse.

Cool pale restlessness — Her skin is cool and pale and she is restless while her heart races to make up the difference for an hour.

Silent urine and confusion — Blood pressure drops to 80/50, urine output stops, and she becomes confused. By the time her blood pressure falls she has already lost over thirty percent of her circulating volume — hypotension is a late sign, never an early indicator.

Two large-bore lines, blood and source control — Two large-bore IVs, warmed isotonic crystalloid, type-specific blood and surgery for source control. Secure the lines for rapid warmed fluid and blood replacement before ever reaching for vasopressors — you cannot squeeze fluid from an empty vascular tank.

14. Bleeding Bea • Warning signs

Hypovolemic Shock



Warning signs

- Heart rate 124
- Respirations 26
- Cool, pale skin
- Restlessness
- Narrowing pulse pressure
- Blood pressure 80/50 (late)
- Urine output stopped
- Confusion

Priority action

Two large-bore IVs, isotonic crystalloid, blood products, and control the source — source control, not passive observation.

14. Bleeding Bea • Check yourself

Hypovolemic Shock



Practice question

Bea's vital signs are HR 124, RR 26, BP 112/88, skin cool and pale. The nurse recognizes this as:

Cover the answer, say yours out loud, then check yourself against the priority action on the previous page and the memory line below.

Remember it

Remember Bleeding Bea whenever you picture an empty fuel tank warning light flashing on a speeding dashboard: her heart races furiously long before the engine finally stalls. Her name reminds you that occult bleeding drops intravascular volume, demanding immediate source control rather than passive observation. For your NCLEX exam, remember that hypotension is never an early indicator; by the time blood pressure falls below normal thresholds, the client has already lost over thirty percent of their circulating capacity. Focus on persistent tachycardia, tachypnea, and a narrowing pulse pressure as the earliest reliable heralds of hypovolemic collapse. Always secure two large-bore intravenous lines for rapid warmed fluid and blood replacement before ever reaching for vasopressors, because you cannot squeeze fluid from an empty vascular tank.

MEMORY LINE

Tank is empty: tachycardia first, hypotension late

15. Anaphylactic Ana

Anaphylaxis



Airway closing — epinephrine IM, thigh, now

Ana gets ten minutes into her first dose of IV penicillin when her lips start to swell and her voice changes. The whole story happens in the time it takes to walk down a hallway.

WHY THIS CHARACTER MATTERS

Ana's immune system overreacts within minutes of an antibiotic dose: hives, swelling lips, stridor, wheezing and a plummeting blood pressure. Epinephrine intramuscularly into the vastus lateralis comes first — antihistamines and steroids are adjuncts, never the first move.

The story arc

1. The setup

A first dose of IV penicillin starts infusing; Ana mentions she has never taken it before.

2. The turn

She scratches at hives on her chest, then says her tongue feels thick.

3. The crisis

Stridor, wheezing, blood pressure 82/40, heart rate 130.

4. The save

The infusion is stopped, IM epinephrine goes into her thigh, oxygen and fluids follow, then diphenhydramine and methylprednisolone.

15. Anaphylactic Ana • What goes wrong

Anaphylaxis



What goes wrong

Ana's immune system overreacts within minutes of an antibiotic dose: hives, swelling lips, stridor, wheezing and a plummeting blood pressure. The whole story happens in the time it takes to walk down a hallway.

NORMAL — THE BASELINE

A first dose infuses without reaction — no hives, no swelling, airway and blood pressure steady.

Closer look

Airway — Ten minutes into her first dose of IV penicillin, her lips start to swell and her voice changes; she says her tongue feels thick — fatal laryngeal edema is the threat.

Breathing — Stridor and wheezing as her immune system overreacts within minutes of the antibiotic dose.

Circulation — A plummeting blood pressure — 82/40 with heart rate 130 — hives on her chest, swelling lips, stridor, and wheezing.

Epinephrine site/time — Epinephrine intramuscularly into the vastus lateralis comes first; antihistamines and steroids are adjuncts, never the first move.

15. Anaphylactic Ana • Warning signs

Anaphylaxis



Warning signs

- Hives — scratching at hives on her chest
- Lips starting to swell; tongue feels thick; voice changes
- Stridor and wheezing
- Blood pressure 82/40, heart rate 130

Priority action

Stop the causative infusion first, then administer IM epinephrine into the thigh immediately; oxygen and fluids follow, then diphenhydramine and methylprednisolone as secondary adjuncts.

15. Anaphylactic Ana • Check yourself

Anaphylaxis



Practice question

A client discharged after an anaphylactic reaction is taught to use an epinephrine auto-injector. Which statement indicates correct understanding?

- A. "I will take an antihistamine first and use epinephrine only if it does not work."
- B. "I will inject epinephrine into my outer thigh first, then call 911 and go to the emergency department even if I feel better."
- C. "If I feel better after the injection, I do not need emergency follow-up."
- D. "I will inject the epinephrine into my buttock and rest at home."

ANSWER: B

Epinephrine IM into the vastus lateralis (outer thigh) is always the first treatment for anaphylaxis, and emergency evaluation is still required afterward because symptoms can recur in a biphasic reaction. Antihistamines are only adjuncts and must never delay epinephrine, which is the trap in allergic-reaction teaching.

Remember it


To ace your exam, remember Anaphylactic Ana as the track coach who always runs straight for the thigh when the airway closes. Her name reminds you that acute anaphylaxis strikes in an instant, demanding intramuscular epinephrine into the vastus lateralis before you touch any other medication vial. Picture Ana sprinting with an auto-injector pointed at her leg to permanently cement the thigh injection site and rapid timing in your memory. On the NCLEX, examiners frequently try to trick you by presenting diphenhydramine or methylprednisolone as appealing first choices for allergic reactions. Your winning clinical rule is absolute: always stop the causative infusion first, then administer intramuscular epinephrine immediately. Antihistamines and corticosteroids are merely secondary adjuncts that will never reverse fatal laryngeal edema or distributive shock in time.

MEMORY LINE

Airway closing — epinephrine IM, thigh, now

16. Embolic Emma

Pulmonary Embolism



Sudden dyspnea — High-Fowler's + oxygen first

Emma sat still on a long flight while a clot packed its bags in her calf. Somewhere over the ocean it boarded the vena cava, flew to her lung and shut down the runway.

WHY THIS CHARACTER MATTERS

Emma's dyspnea hits out of nowhere with pleuritic chest pain, tachycardia and anxiety. First nursing action is High-Fowler's plus oxygen, then notify the provider and prepare for anticoagulation. Watch for hypoxia that doesn't improve and sudden hypotension.

The story arc

1. The setup

A twelve-hour flight, oral contraceptives and a swollen left calf she blamed on her shoes.

2. The turn

Two days later she stands up and cannot breathe. Sharp pain stabs with each inhale.

3. The crisis

SpO₂ 86% on 4 L, heart rate 128, and she says she feels like something terrible is about to happen.

4. The save

High-Fowler's and oxygen first, provider notified, CT angiogram confirms, heparin infusion started with aPTT monitoring.

16. Embolic Emma • What goes wrong

Pulmonary Embolism



What goes wrong

Twelve sedentary hours in a cramped window seat, combined with estrogen's hypercoagulable effect, form a thrombus in her left calf vein. When it breaks free it barrels through the right side of her heart and shuts down the pulmonary runway, creating an acute ventilation-perfusion mismatch.

NORMAL — THE BASELINE

Emma lives out of a carry-on, racing between time zones, and her circulation keeps pace — blood flows from her legs through her heart to her lungs without obstruction. Movement during long travel keeps venous blood from pooling.

Closer look

Calf — Emma's clot assembles silently in the deep veins of her left soleus during a twelve-hour flight: venous stasis from immobility plus hypercoagulability from oral contraceptives — Virchow's triad. She dismisses the mild unilateral calf edema and dull aching as stiff shoes.

Vena cava path — Two days later a shift in position dislodges the clot, and the embolus travels upstream through the inferior vena cava, right atrium, and right ventricle without pause. Never massage the swollen calf — it can release further emboli.

Pulmonary artery — The embolus lodges in the pulmonary arterial tree, occluding blood flow to the alveoli — ventilated but entirely unperfused lung. CT pulmonary angiography confirms bilateral arterial filling defects.

Oxygen field — Her hypoxia is stubborn: SpO₂ stalls at 86% even on four liters of oxygen. Watch for hypoxia that does not improve and sudden hypotension — the red flags of right ventricular failure.

16. Embolic Emma • Warning signs

Pulmonary Embolism



Warning signs

- Sudden-onset dyspnea, striking without warning
- Sharp, knife-like pleuritic chest pain with each inhale
- Respiratory rate 34 breaths per minute
- Sinus tachycardia at 128 beats per minute
- SpO₂ 86% on 4 L of oxygen — hypoxia that does not improve
- An overwhelming sense of impending doom
- pH 7.50, PaCO₂ 28 mm Hg, PaO₂ 56 mm Hg
- D-dimer surged past 1,800 ng/mL
- Unilateral left calf edema and dull aching after the flight

Priority action

High-Fowler's plus oxygen first — elevate Emma into high-Fowler position and give high-flow oxygen by nonrebreather mask before paging the provider, drawing a blood gas, or sending her to CT. Then notify the provider, establish wide-bore IV access, and once CT confirms the obstruction, start the IV heparin bolus and weight-based infusion titrated to an aPTT target of sixty to eighty seconds.

16. Embolic Emma • Check yourself

Pulmonary Embolism



Practice question

A client is being monitored after a confirmed pulmonary embolism and the start of anticoagulation. Which new finding requires the nurse's immediate intervention?

- A. Sharp chest pain that worsens with a deep breath
- B. A feeling of anxiety and restlessness
- C. A respiratory rate of 22 breaths per minute
- D. Blood pressure of 82/50 mmHg with a rapid, weak pulse

ANSWER: D

Sudden hypotension with tachycardia in pulmonary embolism signals obstructive shock as the embolus blocks pulmonary blood flow and the right heart fails, which is immediately life-threatening. Pleuritic pain, anxiety, and mild tachypnea are expected features of the embolism itself and are already being treated. The blood pressure change represents progression and must be escalated at once.

Remember it

To remember Embolic Emma on exam day, envision her transatlantic flight where a packed clot takes off from the calf and shuts down the lung's runway. Her very name reminds you that an embolus is a mobile clot that causes sudden-onset dyspnea, pleuritic pain, and an unshakable feeling that something terrible is about to happen. Notice how her hypoxemia remains stubborn despite baseline oxygen therapy. For the NCLEX, always remember the absolute priority sequence: when a patient presents with sudden respiratory distress, your first independent nursing action is always high-Fowler positioning combined with immediate supplemental oxygen, performed even before you page the provider, obtain an arterial blood gas, or roll the client away to radiology for a computed tomography scan.

MEMORY LINE

Sudden dyspnea — High-Fowler's + oxygen first

17. Rigid Rita

ARDS



Refractory hypoxemia, stiff lungs, white-out on x-ray

Rita survives the sepsis that started it all, then her lungs turn to concrete. The x-ray goes white from edge to edge and every liter of oxygen she gets seems to disappear.

WHY THIS CHARACTER MATTERS

Rita's lungs stiffen and flood 24–48 hours after a major insult. The hallmark is refractory hypoxemia — the PaO_2 will not rise no matter how much oxygen you give. Low tidal volume ventilation, PEEP and prone positioning are the answers.

The story arc

1. The setup

Rita is recovering from severe pancreatitis when her respiratory rate creeps up.

2. The turn

Within 24 hours she needs a non-rebreather, and her SpO_2 still sits at 85%.

3. The crisis

Chest x-ray shows bilateral white-out infiltrates with no heart failure to explain it. Her lungs are stiff and hard to bag.

4. The save

She is intubated on low tidal volumes (6 mL/kg) with high PEEP, sedated, and turned prone for 16 hours a day.

17. Rigid Rita • What goes wrong

ARDS



What goes wrong

Two days into recovery from severe pancreatitis, systemic inflammation leaks into her lungs; within the rapid twenty-four to forty-eight hour window, mild tachypnea becomes full ARDS. Every liter of oxygen seems to disappear — the hallmark is refractory hypoxemia: the PaO₂ will not rise no matter how much oxygen you give.

NORMAL — THE BASELINE

Rita's lungs are spongy and compliant, her alveoli dry and elastic, so air moves easily and oxygen crosses into the blood without effort. Gas exchange keeps pace because the alveolar-capillary membrane stays intact.

Closer look

Alveolar membrane — After a catastrophic systemic insult, inflammatory cytokines damage the alveolar-capillary membrane and permeability surges. Massive amounts of protein-rich fluid pour into the interstitial spaces and alveoli, and surfactant is inactivated and diluted away.

Fluid field — Her chest x-ray shows bilateral white-out infiltrates — flooded alveoli from edge to edge — while a normal BNP excludes heart failure as the cause. This is non-cardiogenic pulmonary edema: the lungs flood even though the heart is not the problem.

Lung recoil — Widespread microatelectasis stiffens the elastic lung tissue until her spongy, compliant lungs turn to rigid, wet concrete. The manual resuscitator bag feels impossibly hard to squeeze against her stiffened airways.

Ventilator waveform — The ventilator is set to lung-protective ventilation: tidal volumes of 6 mL/kg with PEEP at fourteen centimeters of water to recruit collapsed alveoli. Watch the high PEEP closely — sudden hypotension with absent unilateral breath sounds warns of a tension pneumothorax.

17. Rigid Rita • Warning signs

ARDS



Warning signs

- Respiratory rate creeping from eighteen to twenty-eight breaths per minute
- Persistent air hunger and restlessness
- SpO₂ still 85% on a non-rebreather — sliding despite escalating oxygen
- Severe dyspnea with sternocleidomastoid accessory muscle use
- Diffuse fine crackles throughout all lobes
- SpO₂ fixed at eighty-two percent on one hundred percent FiO₂
- PaO₂ 52 mmHg, PaCO₂ 56 mmHg, pH 7.26
- PaO₂ to FiO₂ ratio of 52 — far below the threshold of 100 for severe ARDS
- Bilateral white-out infiltrates on chest x-ray

Priority action

Intubate and ventilate with low tidal volumes (6 mL/kg) and high PEEP to prop open fluid-logged alveoli, with sedation, analgesia, and neuromuscular blockade to eliminate patient-ventilator dyssynchrony. Turn her prone for 16 hours a day to improve ventilation-perfusion matching, and strictly avoid aggressive IV fluids, which would worsen the alveolar flooding.

17. Rigid Rita • Check yourself

ARDS



Practice question

A ventilated client with ARDS is receiving high levels of PEEP. The nurse suddenly notes a drop in blood pressure, absent breath sounds on one side, and worsening oxygen saturation. What complication should the nurse suspect and act on immediately?

- A. Tension pneumothorax related to high airway pressures
- B. Expected improvement from prone positioning
- C. Fluid overload from IV resuscitation
- D. A mucus plug that can be managed at the next routine suctioning

ANSWER: A

High PEEP in stiff, noncompliant ARDS lungs increases the risk of barotrauma, and sudden hypotension with unilateral absent breath sounds is the classic picture of tension pneumothorax, which requires immediate provider notification and emergency intervention. It is not an expected effect of positioning or routine care. Waiting for scheduled suctioning would allow obstructive shock to progress.

Remember it

To make this stick, connect Rigid Rita's nickname to her concrete-stiff lungs that lose compliance and her chest x-ray that displays a dense white-out of flooded alveoli. Remember that her hallmark symptom is refractory hypoxemia, meaning that no matter how much oxygen you deliver, her arterial oxygenation refuses to budge. When preparing for the NCLEX, you must prioritize low tidal volume ventilation at six milliliters per kilogram to prevent fatal volutrauma, apply PEEP to prop open fluid-logged alveoli, and utilize prone positioning to optimize ventilation-perfusion matching. Your crucial NCLEX safety priority is monitoring the high PEEP settings closely; a sudden drop in blood pressure accompanied by absent unilateral breath sounds warns you that a life-threatening tension pneumothorax has developed.

MEMORY LINE

Refractory hypoxemia, stiff lungs, white-out on x-ray

18. Wheezing Wanda

Asthma Exacerbation



A silent chest is worse than a loud one

Wanda's chest whistles all the way to triage. Then, halfway through her second treatment, the whistling stops — and that silence is the most dangerous sound in the room.

WHY THIS CHARACTER MATTERS

Wanda's airways narrow and swell. Expiratory wheeze, prolonged expiration and a falling peak flow. Short-acting albuterol first, then steroids. If the wheezing stops but she is worse, air is not moving at all — that is a respiratory arrest waiting to happen.

The story arc

1. The setup

Cat dander at a friend's house, plus a rescue inhaler she has been using six times a day for a week.

2. The turn

Loud expiratory wheeze, prolonged expiration, peak flow at 45% of her personal best.

3. The crisis

After two treatments the wheezing fades — but she is drowsy, cannot speak in words and her SpO₂ is 88%.

4. The save

Continuous nebulized albuterol with ipratropium, IV corticosteroids, magnesium, and intubation readiness.

18. Wheezing Wanda • What goes wrong

Asthma Exacerbation



What goes wrong

Cat dander at a friend's house triggers the allergic cascade; instead of seeking care she leans on her rescue inhaler six times a day for a week while airway inflammation keeps building. By triage she is in a tripod position, tachypneic and tachycardic, managing only one-word gasps — and then the wheeze disappears because air is no longer moving at all.

NORMAL — THE BASELINE

Wanda's airways are open and she breathes without resistance between rehearsals. Her peak flow sits at her personal best, and her rescue inhaler is a backup, not a daily habit.

Closer look

Bronchial ring — Allergens ignite Wanda's hyperreactive airways: smooth muscle bronchospasm constricts the bronchi while mucosal edema and thick secretions clog the lumen. Forcing air through the narrowed airways generates her high-pitched expiratory wheeze with prolonged expiration.

Mucus layer — An IgE-mediated cascade unleashes histamine, leukotrienes, and prostaglandins across her pulmonary mucosa, and goblet cell secretions thicken inside already narrowing airways. Chest tightness builds like a tightening corset.

Airflow meter — Her peak flow meter tracks the decline: sliding into the yellow caution zone, then to 45% of her personal best — the red zone below 50% that demands rapid rescue treatment. Monitor sequential peak flow responses to every intervention.

Inhaler — Her blue rescue inhaler is the warning light: puffing albuterol six times a day for an entire week signals dangerously failing outpatient control. Short-acting albuterol goes first in the emergency, then steroids — and sedatives are never given to an asthmatic struggling for air.

18. Wheezing Wanda • Warning signs

Asthma Exacerbation



Warning signs

- Rescue inhaler used six times a day for a week
- Nocturnal dry hacking cough and progressive chest tightness
- Loud expiratory wheeze with prolonged expiration
- Peak flow at 45% of personal best — red zone below 50%
- Tripod position with accessory muscle use, speaking only one-word gasps
- SpO₂ 88% on room air
- PaCO₂ of 40 mm Hg — deceptively normal in a hyperventilating asthmatic, signaling muscle exhaustion
- Wheezing suddenly stops — silent chest
- Drowsiness, drooping eyes

Priority action

Sit Wanda in high-Fowler's and give continuous nebulized albuterol with ipratropium first for rapid bronchodilation, with humidified oxygen titrated to keep saturations above 92%. Follow immediately with IV corticosteroids (methylprednisolone) and IV magnesium sulfate. Never give sedatives. Recognize the silent chest as impending respiratory arrest — summon the rapid response team and keep intubation equipment ready at the bedside.

18. Wheezing Wanda • Check yourself

Asthma Exacerbation



Practice question

A client arrives with an acute asthma exacerbation: expiratory wheezing, prolonged expiration, and a peak flow in the red zone. Orders include an albuterol nebulizer treatment, IV methylprednisolone, and oxygen. Which medication should the nurse administer FIRST?

- A. Albuterol by nebulizer
- B. IV methylprednisolone
- C. A sedative to reduce the client's anxiety
- D. An oral antihistamine

ANSWER: A

Inhaled short-acting beta-agonists such as albuterol are the first-line rescue treatment because they rapidly relieve bronchospasm; systemic corticosteroids are given next but take longer to reduce airway inflammation. Sedatives are dangerous in a client struggling to breathe because they can depress respiratory drive. Delaying the bronchodilator to give slower-acting drugs first prolongs hypoxemia.

Remember it

Remember Wheezing Wanda to master the critical trajectory of acute asthma exacerbations on test day. Link her name directly to her prolonged expiratory whistling, but remember her most dangerous symptom: the sudden silence that arrives when air stops moving entirely. Visualize her rescue inhaler emptied six times daily as your classic warning sign of impending decompensation, and picture her falling red-zone peak flow below 50% prompting rapid rescue nebulizers. For your NCLEX exam, remember that a silent chest paired with drowsiness or a normal-looking PaCO₂ in a severely tachypneic asthmatic is not improvement; it is an impending respiratory arrest requiring emergent intervention. Always give inhaled short-acting beta agonists first, follow closely with systemic steroids, and never give sedatives to an asthmatic struggling for air.

MEMORY LINE

A silent chest is worse than a loud one

Meet the 158 Recall Characters

Every character in the binder — see them all at mnemonurse.com/recall-binder



Acidotic Andy

Metabolic Acidosis
Kussmaul breathing, fruity
breath, pH ↓ · HCO_3^- ↓



Alkalotic Alice

Metabolic Alkalosis
Vomiting and diuretics,
tingling fingers, pH ↑ ·
 HCO_3^- ↑



Retaining Ray

Respiratory Acidosis
COPD, CO_2 narcosis, pH ↓
· PaCO_2 ↑



Panicking Pam

Respiratory Alkalosis
Hyperventilation, numb
lips, pH ↑ · PaCO_2 ↓



Potassium Pete

Hyperkalemia
Peaked T waves, muscle
cramps, hold the bananas



Low-K Lena

Hypokalemia
Flat T waves, U waves,
silent belly, never IV push

Rapid-Recall Flashcard Deck

659 cards from every character — cover the answer and say it out loud.

Acidotic Andy

Acidotic Andy — say the memory line from memory.

Kussmaul breathing, fruity breath, pH ↓ ·
HCO₃⁻ ↓

Acidotic Andy

Name a warning sign for Acidotic Andy (Metabolic Acidosis).

pH 7.12 — below 7.35

Acidotic Andy

What goes wrong in Metabolic Acidosis?

Two days of missed insulin plus sugary sodas starve his cells while sugar piles up outside them; osmotic diuresis drains water and electrolytes, and ketone acids overwhelm the buffers into high anion gap metabolic acidosis — his lungs blow off carbon dioxide in compensation, but they cannot outpace the acid.

Acidotic Andy

What is the priority action for Acidotic Andy?

Treat the cause first — insulin plus fluids for DKA, not reflexive bicarb.

Alkalotic Alice

Alkalotic Alice — say the memory line from memory.

Vomiting and diuretics, tingling fingers, pH ↑ ·
HCO₃⁻ ↑

Alkalotic Alice

Name a warning sign for Alkalotic Alice (Metabolic Alkalosis).

Fingers and lips tingling

Alkalotic Alice

What goes wrong in Metabolic Alkalosis?

Three days of vomiting with no fluids kept down, and a furosemide dose she never skipped, drain hydrochloric acid, potassium, and chloride.

Alkalotic Alice

What is the priority action for Alkalotic Alice?

Give normal saline with potassium chloride to replace the chloride she vomited away — isotonic sodium chloride fluid replacement to restore chloride is the priority treatment — and hold the diuretic.

Retaining Ray

Retaining Ray — say the memory line from memory.

COPD, CO₂ narcosis, pH ↓ · PaCO₂ ↑

Retaining Ray

Name a warning sign for Retaining Ray (Respiratory Acidosis).

PaO₂ 60 millimeters of mercury

Your Study Planner

158 characters, 3 a day. Tick each box, then re-test yourself the next morning.

Pace options: Steady = 2 characters a day, Standard = 3 a day, Cram = 6 a day. The grid below is set to the Standard pace. Track the same plan online at mnemonurse.com/recall-binder/planner.

- Day 1**
- ☐ Acidotic Andy - Metabolic Acidosis
 - ☐ Alkalotic Alice - Metabolic Alkalosis
 - ☐ Retaining Ray - Respiratory Acidosis

- Day 2**
- ☐ Panicking Pam - Respiratory Alkalosis
 - ☐ Potassium Pete - Hyperkalemia
 - ☐ Low-K Lena - Hypokalemia

- Day 3**
- ☐ Salty Sal - Hypernatremia
 - ☐ Diluted Dana - Hyponatremia
 - ☐ Chvostek Chuck - Hypocalcemia

- Day 4**
- ☐ Stony Stella - Hypercalcemia
 - ☐ Magnesium Maggie - Magnesium Imbalance
 - ☐ Septic Sam - Distributive Shock

- Day 5**
- ☐ Pumpless Paul - Cardiogenic Shock
 - ☐ Bleeding Bea - Hypovolemic Shock
 - ☐ Anaphylactic Ana - Anaphylaxis

- Day 6**
- ☐ Embolic Emma - Pulmonary Embolism
 - ☐ Rigid Rita - ARDS
 - ☐ Wheezing Wanda - Asthma Exacerbation

- Day 7**
- ☐ Consolidated Carl - Pneumonia
 - ☐ Vented Vince - Mechanical Ventilation
 - ☐ Ketone Kyle - Diabetic Ketoacidosis